

A single domain of the ZP2 zona pellucida protein mediates gamete recognition in mice and humans

Matteo A. Avella, Boris Baibakov, and Jurrien Dean

Laboratory of Cellular and Developmental Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892

The extracellular zona pellucida surrounds ovulated eggs and mediates gamete recognition that is essential for mammalian fertilization. Zonae matrices contain three (mouse) or four (human) glycoproteins (ZP1–4), but which protein binds sperm remains controversial. A defining characteristic of an essential zona ligand is sterility after genetic ablation. We have established transgenic mice expressing human ZP4 that form zonae pellucidae in the absence of mouse or human ZP2. Neither mouse nor human sperm bound to these ovulated eggs, and these female mice were sterile after in vivo

insemination or natural mating. The same phenotype was observed with truncated ZP2 that lacks a restricted domain within ZP2^{51–149}. Chimeric human/mouse ZP2 isoforms expressed in transgenic mice and recombinant peptide bead assays confirmed that this region accounts for the taxon specificity observed in human–mouse gamete recognition. These observations in transgenic mice document that the ZP2^{51–149} sperm-binding domain is necessary for human and mouse gamete recognition and penetration through the zona pellucida.

Introduction

Monospermic fertilization is required for successful development in mice and humans. However, the molecular basis of sperm–egg interaction remains incompletely understood despite decades of investigation. Ovulated eggs are surrounded by an extracellular zona pellucida, to which sperm bind and penetrate before gamete fusion. After fertilization, the zona matrix is modified so that sperm do not bind to the early embryo. Humans have four genetic loci encoding ZP1, ZP2, ZP3, and ZP4 (Spargo and Hope, 2003), but mouse *Zp4* is a pseudogene (Lefièvre et al., 2004), and the mouse zona pellucida contains only three glycoproteins (Bleil and Wassarman, 1980). Given the simple structure of the zona pellucida, it has been surprisingly difficult to genetically define a zona protein that is essential for fertilization and the postfertilization block to polyspermy.

To identify the zona ligand for sperm recognition, we have exploited two physiological dichotomies. One is that sperm bind to the zona pellucida surrounding eggs before, but not after, fertilization. The only biochemically documented change in the zona matrix is the postfertilization cleavage of ZP2 (Bleil et al., 1981; Bauskin et al., 1999), and the site has been defined as ¹⁶⁶LA↓DE¹⁶⁹ in mice (Gahlay et al., 2010). If mutated in transgenic mice or if the cleaving enzyme, ovastacin, is genetically

ablated, ZP2 remains intact and sperm bind de novo to the zona pellucida surrounding early embryos despite fertilization (Gahlay et al., 2010; Burkart et al., 2012). The second dichotomy is that human sperm bind to human, but not mouse, zonae pellucidae (Bedford, 1977).

To capitalize on these differences, we have established transgenic mouse lines lacking ZP2 and expressing truncated or human/mouse chimeric isoforms to define ZP2^{51–149} as the zona ligand for human and mouse sperm recognition, and document that in its absence female mice are sterile.

Results and discussion

Female mice that form a zona pellucida lacking ZP2 are sterile


Formation of the extracellular zona matrix is mediated by zona domains (260 aa motifs with conserved cysteine residues) present near the C terminus of each secreted zona protein (Bork and Sander, 1992). The human zona pellucida is composed of four (ZP1–4) and the mouse of three (ZP1–3) glycoproteins (Bleil and Wassarman, 1980; Bauskin et al., 1999). Mouse ZP1, the least

Correspondence to Jurrien Dean: jurriend@helix.nih.gov

Abbreviations used in this paper: BAC, bacterial artificial chromosome; HTF, human tubal fluid.

This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

Figure 1. A zona pellucida is formed with ZP4 in the absence of ZP2, and female mice are sterile. (A) Phylogeny of mouse and human zona proteins indicate two clades; one composed of ZP1, ZP2, and ZP4 and the other of ZP3. There is no mouse ZP4 protein because of multiple stop and missense codons in the cognate gene. Mya, million years ago. (B) Schematic representation of the four zona pellucida proteins with 8 or 10 conserved cysteine residues. The resultant disulfide bonds differ in the zona domains of the ZP1/2/4 and ZP3 clades and are indicated as A and B, respectively. The postfertilization cleavage site is marked on ZP2, and both ZP1 and ZP4 contain trefoil domains. (C) Glutaraldehyde-fixed, plastic-embedded ovarian sections (3 μ m) from 8–10-wk-old normal, *Zp2^{Null}*, *moQuad^(huZP4)*, and *moQuad-Zp2^{Null}* mice were stained with periodic acid Schiff's reagent to highlight the zona pellucida (arrows) and counterstained with hematoxylin. (D) Formaldehyde-fixed *moQuad-Zp2^{Null}* eggs stained with protein-specific monoclonal antibodies. Fluorescent and DIC images were merged and faux colored. (E) Mouse sperm binding to *moQuad^(huZP4)* and *moQuad-Zp2^{Null}* eggs. Inset, 2.5 \times magnification. *Zp3^{EGFP}* mouse eggs (green zona) and mouse two-cell embryos were positive and negative sperm-binding controls, respectively. Schematics to the left reflect protein composition of the zonae pellucidae with the source of sperm below.



abundant protein, is not required for formation of the zona pellucida or fertility (Rankin et al., 1999). In the absence of ZP2, a thin zona matrix is formed around growing oocytes that does not persist in ovulated eggs (Rankin et al., 2001), and no zona matrix is formed in the absence of ZP3 (Liu et al., 1996; Rankin et al., 1996). Based on the phylogeny (Larkin et al., 2007) of human and mouse zona proteins, ZP1/ZP2/ZP4 fall into one clade and ZP3 into another (Fig. 1 A). The zona domains of ZP2 and ZP3 have 10 and 8 conserved cysteine residues, respectively, the linkage of which differs in the two clades (Fig. 1 B; Boja et al., 2003). We reasoned that if members of the first clade could substitute for one another, the presence of ZP4 along with ZP1 might permit formation of a matrix in the absence of mouse ZP2 and provide a loss-of-function assay for sperm–egg recognition. Because mouse *Zp4* is a pseudogene (Lefèvre et al., 2004), human *ZP4* was expressed in transgenic mice (Yauger et al., 2011) to establish a mouse line designated *moQuad* (*moZP1*, *moZP2*, *moZP3*, and *huZP4*). After the appropriate crosses, mice lacking mouse ZP2 in the presence of ZP4 were established and designated *moQuad-Zp2^{Null}* (*moZP1*, *moZP3*, and *huZP4*; Table S1).

In the presence of *huZP4*, *moQuad^(huZP4)* (ZP1,2,3,4) and *moQuad-Zp2^{Null}* (ZP1,3,4) transgenic lines form zonae pellucidae during oocyte growth that is similar to normal oocytes (Fig. 1 C and Table S1). The zona pellucida persists after eggs are ovulated into the oviduct, and the composition of the *moQuad-Zp2^{Null}* zona matrix was confirmed with monoclonal antibodies that documented the absence of mouse ZP2 (Fig. 1 D). Ovulated eggs in cumulus (hyaluron interspersed with follicular cells) from *moQuad^(huZP4)* and *moQuad-Zp2^{Null}* mice were inseminated

with mouse sperm using *Zp3^{EGFP}* mouse eggs (green zona) and normal mouse two-cell embryos, respectively, as positive and negative controls. Mouse sperm bound to *moQuad^(huZP4)* (41.4 ± 2.5 , $n = 30$), but not to *moQuad-Zp2^{Null}* eggs (1.6 ± 0.3 , $n = 43$; Fig. 1 E), and the latter, but not the former, mouse line was sterile (Table S2). Thus, mouse ZP2 is required for in vitro sperm binding and in vivo mouse fertility.

Truncated ZP2 does not support mouse sperm binding

The secreted ZP2 ectodomain (35–633 aa) lacks the signal peptide (1–34) that directs ZP2 into the endosomal pathway and the C terminus (634–713) that includes a transmembrane domain. After fertilization, the ectodomain is cleaved near the N terminus ($^{166}\text{LA}\downarrow\text{DE}^{169}$) by ovastacin, an egg cortical granule metalloendoprotease, after which sperm no longer bind to the zona pellucida (Gahlay et al., 2010; Burkart et al., 2012). To investigate the importance of the N terminus in mouse gamete recognition, DNA recombineering was used to construct a transgene lacking ZP2^{51–149} (*moZp2^{Trunc}*; Fig. 2 A and Fig. S1 A). After crossing into the *Zp2^{Null}* background, female mice reconstituted a zona pellucida that was thinner than normal. Therefore, these lines were crossed with *huZP4* transgenic mice to establish *moQuad-Zp2^{Trunc}* mice that developed a more robust zona matrix during oocyte growth (Fig. 2 B).

The zona pellucida surrounding ovulated eggs from *moQuad* and *moQuad-Zp2^{Trunc}* mice were analyzed by confocal microscopy using ZP2 domain-specific monoclonal antibodies. The zonae from both genotypes reacted with antibodies to ZP1, ZP2^{C-term}, ZP3, and *huZP4*. However, the monoclonal antibody




Figure 2. Truncated ZP2 does not support sperm binding, and female mice are sterile. (A) Representation of secreted ectodomains of normal mouse ZP2^{35–633} and truncated ZP2 lacking ZP2^{51–149}. Cysteine residues, yellow. Monoclonal antibodies that bind N and C terminal to the postfertilization cleavage site (arrowhead) and zona domains are indicated above. (B) Ovarian histology of moZp2^{Trunc} and moQuad-Zp2^{Trunc} transgenic mice in Zp2^{Null} background as in Fig. 1 C. (C) moQuad^(huZP4) and moQuad-Zp2^{Trunc} eggs stained with domain-specific monoclonal antibodies as in Fig. 1 D. (D) Immunoblot of eggs (15) from moQuad^(huZP4) (1) and moQuad-Zp2^{Trunc} (2) mice stained with domain-specific monoclonal antibodies. Molecular masses are indicated on the left. (E) Mouse sperm binding to Zp2^{Trunc} and moQuad-Zp2^{Trunc} eggs as in Fig. 1 E.

to the N terminus of ZP2 did not react with the zona pellucidae surrounding moQuad-Zp2^{Trunc} eggs, although it recognized normal ZP2 in the moQuad (Fig. 2 C). On immunoblots, the monoclonal antibody to the ZP2^{N-term} detected a 120-kD band in the zona pellucida isolated from moQuad, but not moQuad-Zp2^{Trunc} eggs (Fig. 2 D, left). A monoclonal antibody to the ZP2^{C-term} detected a 120-kD band in the zona pellucida isolated from moQuad eggs and a 92-kD band in eggs from moQuad-Zp2^{Trunc} mice (Fig. 2 D, right). These observations are consistent with deletion of 99 amino acids (ZP2^{51–149}) in the N terminus of ZP2 of the zona pellucida surrounding moQuad-Zp2^{Trunc} eggs.

Mouse sperm did not bind to either Zp2^{Trunc} (0.6 ± 0.2 , $n = 22$) or moQuad-Zp2^{Trunc} (2.3 ± 1.3 , $n = 30$) eggs (Fig. 2 E), and, when mated with normal males, moQuad-Zp2^{Trunc} females were sterile (Table S2). Based on these observations in transgenic mice, we conclude that the N terminus of mouse ZP2 is necessary for gamete recognition in vitro and in vivo.

Human sperm recognize human but not mouse ZP2

To extend these observations to human biology, huQuad^(huZP1–4) mice containing all four human proteins and none of the endogenous mouse proteins (Baibakov et al., 2012) were used to establish huQuad-ZP2^{Null} (huZP1,3,4) mouse lines (Table S1). Both lines formed a zona surrounding growing oocytes within the ovary (Fig. 3 A), and, using monoclonal antibodies specific to the human proteins, the absence of huZP2 in huQuad-ZP2^{Null} ovulated eggs was confirmed (Fig. 3 B). After insemination with human semen, sperm bound robustly to huQuad^(huZP1–4) eggs (59.8 ± 5.3 , $n = 37$), but rarely to huQuad-ZP2^{Null} eggs (1.5 ± 0.3 , $n = 36$; Fig. 3 C).

To ascertain if mouse ZP2 exerted a non-taxon-specific effect on the zona structure important for gamete recognition, mouse Zp2 was expressed in huQuad-ZP2^{Null} eggs. However, human sperm did not bind to the zona pellucida surrounding huQuad-ZP2^{Null}; Zp2^{Mo} eggs (0.7 ± 0.2 , $n = 30$; Fig. 3 C).

To determine if these observations pertain in vivo, an artificial insemination assay was established using mouse sperm as a positive control for fertility and for accumulation of sperm in the perivitelline space. After transcervical insemination (Fig. S2 A), control (*Cd9*^{+/–}) mice were fertile, albeit with smaller litters (2.0 ± 0.54 vs. 10.4 ± 0.81 for natural mating). *Cd9*^{Null} eggs to which sperm will not fuse (Le Naour et al., 2000) accumulated 1–6 mouse sperm in their perivitelline spaces (between the inner aspects of the zona matrix and the plasma membrane) in 9 of 113 eggs (Fig. 3 D). Subsequently, huZP2^{Rescue} (huZP2 replaces moZP2) and huZP3^{Rescue} (huZP3 replaces moZP3) mice (Table S1) were inseminated with human sperm (3×10^7) and eggs recovered from the oviduct were examined by confocal microscopy to detect sperm in the perivitelline space where they accumulated, unable to fuse with mouse eggs. Of 29 eggs recovered from four huZP2^{Rescue} mice, human sperm were detected in the perivitelline space of four eggs (one egg per female), which is consistent with the small litters observed in control (*Cd9*^{+/–}) female mice. No sperm were observed in the perivitelline space of 28 huZP3^{Rescue} eggs (Fig. 3 E). To facilitate access to ovulated eggs, human sperm also were transferred directly to the oviduct (Fig. 3 F). After 2 h of incubation in vivo, 1–5 human sperm were observed in the perivitelline space of 22–77 eggs (29%) from huZP2^{Rescue} mice (Fig. 1 G). No sperm were observed in the perivitelline space of 85 control huZP3^{Rescue} eggs, which is




Figure 3. Human sperm binding to the zona pellucida requires human ZP2. (A) Ovarian histology from huQuad(*huZP1-4*) and huQuad-ZP2^{Null} mice as in Fig. 1 C. (B) Eggs from huQuad-ZP2^{Null} stained with monoclonal antibodies as in Fig. 1 D. (C) Human sperm binding to huQuad(*huZP1-4*), huQuad-ZP2^{Null}, and huQuad-ZP2^{Null}; *Zp2*^{Mo} eggs (as in Fig. 1 E) using noninseminated human oocytes and mouse *Zp3*^{EGFP} eggs (green zona) as positive and negative controls, respectively. (D) Litter sizes after transcervical insemination of control (*Cd9*^{+/+}) mice compared with natural mating (top). Sperm in the perivitelline space (PVS) of *Cd9*^{-/-} eggs after transcervical insemination with mouse sperm (bottom). Recovered eggs (left) and the number of sperm in PVS (right). Arrows indicate sperm. (E) As in D (bottom) but with huZP2^{Rescue} (top) and huZP3^{Rescue} (bottom) eggs after transcervical insemination with human sperm. (F) In vivo oviduct transfer of human sperm (2.3×10^3 sperm in $0.5 \mu\text{l}$) to hormonally stimulated, anesthetized huZP2^{Rescue} and huZP3^{Rescue} female mice. (G) As in E, but after in vivo oviductal transfer. (H) Normal mouse, huZP2^{Rescue}, huQuad eggs, and human oocyte stained with antibody to the sialyl-Lewis^x antigen.

consistent with their inability to support human sperm binding in vitro (Baibakov et al., 2012). Thus, human sperm bind and penetrate the zona pellucida of huZP2^{Rescue}, but not huZP3^{Rescue}, eggs in vitro and in vivo.

Mouse sperm, lacking taxon-specific gamete recognition (Bedford, 1977), also bound to huQuad(*huZP1-4*) expressing

human ZP2 (35.3 ± 1.9 , $n = 52$) and the huQuad-ZP2^{Null}; *Zp2*^{Mo} (36.3 ± 2.2 , $n = 36$) expressing mouse ZP2, but not the huQuad-ZP2^{Null} eggs missing both mouse and human ZP2 (2.4 ± 0.3 , $n = 44$; Fig. S2 B). After mating, huQuad(*huZP1-4*) and huQuad-ZP2^{Null}; *Zp2*^{Mo} female mice containing human or mouse ZP2 in their zonae pellucidae were fertile, but huQuad-ZP2^{Null} female




Figure 4. Taxon-specific sperm recognition of the N terminus of chimeric ZP2. (A) Ectodomains of huZP2, chimeric hu/moZP2, and chimeric mo/huZP2 proteins. Red and green, human and mouse protein, respectively. Yellow, conserved cysteine residues. Postfertilization cleavage site (arrowhead) and zona domains are indicated above. (B) Ovarian histology of hu/moZP2^{Rescue} and mo/huZP2^{Rescue} as in Fig. 1 C. (C) huZP2^{Rescue}, hu/moZP2^{Rescue}, and mo/huZP2^{Rescue} eggs stained with domain-specific monoclonal antibodies as in Fig. 2 C. (D) Human sperm binding to huZP2^{Rescue}, hu/moZP2^{Rescue}, and mo/huZP2^{Rescue} eggs as in Fig. 3 C. (E) Schematic of human (red) and mouse (green) recombinant peptides in which mouse ZP2⁵²⁻⁸³, ZP2⁸⁵⁻¹⁰¹, or ZP2¹⁰³⁻¹³³ replace the corresponding human sequence. The green bar under the huZP2 protein was deleted in the mouse ZP2^{Trunc}. Cysteine residues, yellow bars. Predicted N-glycosylation sites, blue bars with asterisks. Arrowhead, di-acidic residues and potential ovastacin cleavage sites. (F) Capacitated human sperm binding to chimeric ZP2 peptide beads. huZP2³⁹⁻¹⁵⁴ and moZP2³⁵⁻¹⁴⁹ peptides were positive and negative controls, respectively. DIC (top) and confocal z projection (bottom) images after staining with Hoechst. (G) Box plots reflect the median (vertical line) number of human sperm binding to peptide beads (left) and data points within the 10th and 90th percentiles (error bars). Boxes include the middle two quartiles and outliers are indicated by dots.

mice lacking mouse and human ZP2 were sterile (Table S2). Collectively, these results indicate that ZP2 is necessary for human and mouse sperm binding and penetration through the zona pellucida, and for mouse fertility.

Earlier models of gamete recognition had focused on the role of specific glycans as ligands for sperm binding (Florman and Wassarman, 1985; Bleil and Wassarman, 1988; Miller et al., 1992; Chen et al., 1998). However, continued fertility after mutation of attachment sites (Liu et al., 1995; Gahlay et al., 2010) and genetic ablation of specific galactosyltransferases (Thall et al., 1995; Lowe and Marth, 2003; Shi et al., 2004; Williams et al., 2007) have not supported the candidacy of any of the proposed glycans. More recently, the sialyl-Lewis^x antigen has been reported to mediate human sperm binding to human zonae pellucidae (Pang et al., 2011). Although the sialyl-Lewis^x antigen was detected in the zona pellucida surrounding control human oocytes, it was not present in the zona matrix formed by normal, huZP2^{Rescue}, or huQuad^(huZP1-4) eggs (Fig. 3 H) and cannot account for the observed binding of human sperm under these experimental conditions.

The ZP2 domain required for sperm binding regulates taxon-specific gamete recognition

The ectodomains of human (602 aa) and mouse (599 aa) ZP2 share 62% amino acid identity, but the N termini of the two ZP2 proteins are only 48% identical. Thus, we questioned whether the taxon specificity is mediated by the gamete recognition domain found to be essential for normal mouse sperm binding and fertility in Zp2^{Trunc} mice. Using DNA recombineering, genomic regions encoding the N termini of moZp2 and huZP2 were replaced with the corresponding human exons encoding huZP2²²⁻¹⁶⁴ and mouse exons encoding moZP2¹⁸⁻¹⁵⁶, respectively (Fig. S1 A). After establishment by pronuclear injection, each transgenic line was crossed into the mouse Zp2^{Null} line to eliminate endogenous ZP2 protein and designated hu/moZP2^{Rescue} and mo/huZP2^{Rescue} mice, respectively (Fig. 4 A). The mice appeared normal and a robust zona pellucida was observed in ovarian sections from hu/moZP2^{Rescue} and mo/huZP2^{Rescue} female mice (Fig. 4 B).

Mouse ZP1 and ZP3 were present in the zona pellucida surrounding ovulated eggs from each of the three transgenic mouse




Figure 5. Model of gamete recognition on the surface of the zona pellucida. The mouse zona pellucida (aquamarine) is composed of ZP1, ZP2, and ZP3, and surrounds ovulated eggs and early embryos. Sperm, capacitated by passage through the female reproductive tract, bind on the surface of the zona pellucida to an N-terminal domain of ZP2 in unfertilized eggs. After sperm acrosome exocytosis and penetration of the zona matrix, gametes fuse at fertilization and activate the egg. This triggers egg cortical granule migration and fusion with the plasma membrane, which releases ovastacin, a zinc metalloendoprotease that cleaves ZP2 at ¹⁶⁶LAJDE¹⁶⁹. The immediate postfertilization block to polyspermy prevents additional sperm from fusing with eggs or penetrating through the zona pellucida matrix. The most definitive block is secondary to the proteolytic destruction of the sperm binding domain at the N terminus of ZP2. If sperm do not bind, they will not penetrate nor fuse with the egg's plasma membrane.

lines (Fig. 4 C). The zona pellucida from huZP2 rescue mice (Table S1) reacted with monoclonal antibodies to huZP2^{N-Term} and not with either mouse-specific monoclonal antibody. The zona pellucida of the chimeric hu/moZP2 eggs reacted with huZP2^{N-term} and moZP2^{C-term}, but not moZP2^{N-term} monoclonal antibodies, and the zona pellucida surrounding the chimeric mo/huZP2 rescue eggs reacted with antibodies to the N terminus, but not the C terminus, of mouse ZP2 (Fig. 4 C). Sperm binding was assessed in vitro using noninseminated human oocytes as a positive control and normal mouse eggs with a green zona (Zp3^{EGFP}) as a negative control. Human sperm bound to the zona pellucida surrounding huZP2 (30.7 ± 1.3 , $n = 52$) and hu/moZp2 (22.5 ± 1.7 , $n = 69$), but not to the zona surrounding mo/huZP2 rescue eggs (2.9 ± 0.5 , $n = 49$; Fig. 4 D). Based on these observations, we conclude that a domain near the N terminus of ZP2 mediates the taxon specificity of human sperm binding to the zona pellucida.

Using ZP2³⁹⁻¹⁵⁴ as a backbone, a recombinant baculovirus encoding N-terminal chimeric human and mouse ZP2 peptides was constructed in which mouse replaced human sequence between cysteine residues to systematically express moZP2⁵²⁻⁸³, moZP2⁸⁵⁻¹⁰¹, and moZP2¹⁰³⁻¹³³ in place of the corresponding human sequence (Fig. 4 E and Fig. S2 C). Peptide beads were incubated with human sperm and washed to remove loosely adherent sperm, and binding was quantified from z projections of confocal images. HuZP2³⁹⁻¹⁵⁴ (15.0 ± 0.9 , $n = 43$) and moZP2³⁵⁻¹⁴⁹ (1.2 ± 0.2 , $n = 43$) peptide beads were used as positive and

negative controls, respectively. Replacement of human sequence with moZP2⁵²⁻⁸³ (3.2 ± 0.2 , $n = 16$) dramatically decreased human sperm binding, whereas chimeric peptides with moZP2⁸⁵⁻¹⁰¹ (10.5 ± 1.0 , $n = 14$) or moZP2¹⁰³⁻¹³³ (11.7 ± 0.8 , $n = 14$) had a more minimal effect (Fig. 4 F). These observations were quantified with box plots (Fig. 4 G), which indicate that the region between the first two cysteine residues (huZP2⁵⁵⁻⁸⁸; moZP2⁵²⁻⁸³) near the N terminus of ZP2 plays a significant role in human gamete recognition.

Perspective

ZP2 was first proposed as a primary sperm-binding ligand in *Xenopus laevis* (Tian et al., 1999) and more recently in humans (Baibakov et al., 2012). However, like all other candidates, it had not been defined as essential by genetic ablation. We now report that transgenic mice expressing human ZP4 form a zona pellucida in the absence of ZP2. Using this model system, we have identified a domain within ZP2 that accounts for the taxon specificity of human sperm binding to the zona pellucida and, by genetic ablation, document that this domain is essential for female mouse fertility. Collectively, these results are consistent with a model (Fig. 5) in which mouse and human sperm bind to ZP2⁵¹⁻¹⁴⁹ and hyperactive sperm penetrate the zona matrix to fuse and fertilize ovulated eggs. This triggers egg cortical granule exocytosis in which ovastacin, a metalloendoprotease (Quesada et al., 2004) encoded by *Astl*, is released and cleaves upstream of a di-acidic motif (¹⁶⁶LAJDE¹⁶⁹) in ZP2 (Burkart et al., 2012). Although a single cleavage site is reported in human ZP2 (Bauskin et al., 1999), multiple degradation products are observed in mouse ZP2 (Burkart et al., 2012). In each case, the postfertilization cleavage of ZP2 provides a definitive block to polyspermy. Sperm that do not bind to the zona pellucida cannot penetrate the zona matrix or fuse with the egg plasma membrane. The destruction of the N terminus of ZP2 provides an effective block to polyspermy and ensures the monospermic fertilization required for the successful onset of development.

Materials and methods

Transgenic mouse lines

Targeted mutagenesis with a PGK-neomycin cassette was used to genetically ablate *Zp1* (replaced exon 1–3), *Zp2* (replaced exon 1–2), *Zp3* (disrupted exon 1), and *Cd9* (replaced promoter and exon 1) in embryonic stem (ES) cells. Correctly targeted ES cells were identified for injection into blastocysts to establish mouse lines with null alleles (Rankin et al., 1996, 1999, 2001; Le Naour et al., 2000). *Cd9* null mice were a gift of C. Boucheix (Institut National de la Santé et de la Recherche Médicale, Villejuif, France). Mouse lines expressing human zona proteins were established after pronuclear injection of genomic DNA encoding human ZP1 (11.9 kb including 2.2 promoter, 8.1 coding region, and 1.5 of 3' flank), ZP2 (16.8 kb including 2.3 promoter, 14 kb coding region, and 0.5 kb of 3' flank), ZP3 (16 kb including 6.0 kb promoter, exons 1–4 followed by cDNA encoding exons 5–8 and a BGH polyadenylation signal), and ZP4 (11.6 kb including 2.4 promoter, 8.2 coding region, and 1.0 of 3' flank; Rankin et al., 1998, 2003; Yauger et al., 2011; Baibakov et al., 2012). In addition, three new mouse lines were established using chimeric hu/moZp2, chimeric mo/huZP2, and moZp2^{trunc} transgenes (Fig. S2 A) constructed by GalK DNA recombineering (Warming et al., 2005).

To establish the new lines, bacterial artificial chromosome (BAC) DNA (Life Technologies) that include either mouse *Zp2* (RP23-6513) or human *ZP2* (RP11-1023A8) were transformed into SW102 bacterial cells containing the λ prophage recombineering system (Liu et al., 2003). For hu/moZp2, human genomic DNA encoding huZP2⁴¹⁻¹⁶⁸ (bp 26 in exon

2 to bp 21 in exon 6) replaced mouse genomic DNA encoding moZP2^{37–165} (bp 59 in exon 2 to bp 6 in exon 5). To construct this transgene, a PCR fragment (1,331 bp) containing the *galK* operon flanked by 50 bp homologous to moZp2 gene 5' and 3' of the sequence encoding moZP2^{22–161} protein was amplified (huZP2-Galk primers; Table S3) using NEB Phusion (New England Biolabs, Inc.). After digestion with DpnI and overnight gel purification (0.7% agarose, 15 V, 16 h), the PCR fragment was electroporated into the BAC containing SW102 cells, and recombinants were selected by growth on minimal media with galactose. Using a clone from this first step, the *galK* cassette was replaced by recombineering with a second PCR fragment (5,783 bp) encoding huZP2^{41–168} protein with 100-bp arms homologous to moZP2 on either side (moZp2-huZP2 primers; Table S3). Mutant clones were selected on minimal media with 2-deoxy-galactose and confirmed by DNA sequencing of PCR products using gene specific primers (Table S3) to isolate 5' and 3' junction fragments.

In a similar recombineering strategy, DNA encoding huZP2^{22–161} protein was replaced with sequence encoding moZP2^{18–156} using huZP2-Galk and huZP2-moZp2 primers (Table S3), and DNA encoding moZP2^{51–149} protein was removed using moZp2-Galk and bridging moZp2 primers (Table S3) to establish the mo/huZP2 and moZp2^{trunc} transgenes, respectively. For the mo/huZP2 transgene, mouse genomic DNA encoding moZP2^{18–156} (bp 1 in exon 2 to bp 150 in exon 5) replaced human genomic DNA encoding huZP2^{22–161} (bp 1 in exon 2 to bp 152 in exon 5), and for the moZp2^{trunc} transgene, DNA encoding moZP2^{51–149} (bp 59 in exon 2 to bp 27 in exon 6) was deleted (Fig. S1 A).

NotI fragments containing the hu/moZp2 (17.9 kb including 2.3 and 0.9 kb of the 5' and 3' flanking regions, respectively), mo/huZP2 (16.8 kb including 1.5 and 1.7 kb of the 5' and 3' flanking regions, respectively), and truncated moZp2^{trunc} (13.9 kb including 2.3 and 0.9 kb of the 5' and 3' flanking regions, respectively) transgenes were retrieved from the BAC with pl253 (Gahlay et al., 2010), and the fidelity of coding regions was confirmed by DNA sequencing. After gel purification, the transgenes were injected into the male pronucleus of fertilized FVB/N eggs by the Taconic Transgenic Mouse Facility. At least two founders were established for each transgene and crossed into mouse Zp1-3 null and human ZP1-4 transgenic mouse lines.

Genotyping

Hu/moZp2 mice were genotyped by PCR [(95°C for 30 s, 58°C for 30 s, 72°C for 1 min) × 30 cycles, 72°C for 7 min, and 4°C for >30 min] using primers in intron 5 of human ZP2 and intron 6 of mouse Zp2, mo/huZP2 mice were genotyped using primers in intron 4 of mouse Zp2 and intron 5 of human ZP2, and moZp2^{trunc} mice were genotyped by PCR using primers in intron 2 and intron 5 of mouse Zp2 (Table S4). Each transgenic line was crossed into the Zp2^{Null} background (Rankin et al., 2001) to establish hu/moZp2, mo/huZP2, and moZp2^{trunc} rescue lines. To avoid ambiguity of PCR product sizes, the normal Zp2 allele was detected by three distinct primer sets for hu/moZp2 (Set 1), mo/huZP2 (Set 2), and moZp2^{trunc} (Set 3) rescue lines (Fig. S1 A; Table S4).

Expression of transgenes

Expression of normal Zp2, as well as hu/moZp2, mo/huZP2, and moZp2^{trunc} transgenes, was detected by RT-PCR using as gene-specific primer sets (Table S5) and total RNA isolated from 8–12-wk-old mouse tissue (Fig. S2 B). *Gapdh* transcripts served as a control for RNA integrity (Baibakov et al., 2012).

Microscopy

Samples were mounted in PBS, and images of eggs, embryos, and beads were obtained with a confocal microscope (LSM 510; Carl Zeiss) using a 63×/1.2 NA water immersion objective lens at room temperature (Baibakov et al., 2007; Yauger et al., 2011). LSM 510 images were exported as full-resolution TIF files and processed in Photoshop CS5.5 (Adobe) to adjust brightness and contrast. Alternatively, confocal optical sections were projected to a single plane with maximum intensity and combined with DIC images of eggs or peptide beads using LSM image software.

Ovarian histology and immunohistochemistry

Mouse ovaries were fixed in glutaraldehyde and embedded in glycol methacrylate before staining with periodic Schiff's acid and hematoxylin (Yauger et al., 2011). Ovulated eggs were fixed in 2% paraformaldehyde before staining with rat or mouse monoclonal antibodies (1:50) specific to: moZP1 (Rankin et al., 1999), N terminus of moZP2 (East and Dean, 1984), C terminus of moZP2 (Rankin et al., 2003), moZP3 (East et al., 1985), huZP1 (Ganguly et al., 2010), huZP2 (Rankin et al., 2003), huZP3 (Rankin et al., 1998), and huZP4 (Bukovsky et al., 2008). Monoclonal antibodies to human

ZP1 and ZP4 were a gift from S. Gupta (National Institute of Immunology, New Delhi, India). Primary antibody binding was detected with goat anti-mouse Alexa Fluor 568 (1:100) or goat anti-rat Cy3 (1:100) antibodies (Life Technologies). Monoclonal antibody CSLEX1 (BD) that binds to the sialylLewis^x antigen was diluted 1:50 and detected with goat anti-mouse IgM-FITC (1:20; Life Technologies). Images were obtained at room temperature on a microscope (Carl Zeiss; Axioplan 2) equipped with a 40×/0.75 NA lens using a camera (AxioCam 1Cc1) and AxioVision software (all from Carl Zeiss).

Assessment of sperm binding and fertility

To assay mouse sperm binding to the zona pellucida surrounding normal and transgenic mouse eggs, sperm were released from cauda epididymides, capacitated in human tubal fluid (HTF; EMD Millipore) supplemented with 0.4% BSA (Sigma-Aldrich) for 40 min (37°C, 90% N₂, 5% O₂, and 5% CO₂), and added to eggs in cumulus and embryos in 100 µl of HTF, 0.4% BSA at a final concentration of 10⁵ ml⁻¹ progressive motile sperm as determined by a HTM-IVOS (Version 12.3) motility analyzer (Hamilton Thorne; Gahlay et al., 2010). Zp3^{EGFP} mouse eggs (green zona) and two-cell embryos served as positive and negative wash controls, respectively. Samples were fixed in 2% paraformaldehyde and stained with Hoechst to identify nuclei. Bound sperm were quantified from z projections obtained by confocal microscopy (Baibakov et al., 2007), and results reflect the mean ± SEM from at least three independently obtained samples each containing 10–50 mouse eggs/embryos.

To assay human sperm binding, individual aliquots (0.5 ml) of liquefied human semen (Genetics & IVF Institute Fairfax Cryobank) were added to an Eppendorf tube (2.0 ml) containing 0.5 ml of 40% of PureSperm (Nidacon) layered over 0.5 ml of 80% PureSperm. After centrifugation (swinging bucket, 20 min × 300 g, 20°C) and removal of the supernatant, sperm were resuspended in the residual buffer and transferred into 1.0 ml HTF. After a second centrifugation (5 min × 300 g), sperm were resuspended in 0.2 ml of HTF/BSA, and 2–4 aliquots were mixed before evaluation by HTM-IVOS. Sperm were then diluted in HTF, 0.5% BSA to 10⁷ ml⁻¹ progressive motile sperm. Transgenic eggs in cumulus, normal eggs, Zp3^{EGFP} eggs (Zhao et al., 2002), and noninseminated, immature human oocytes were incubated in droplets (100 µl) of sperm (10⁷) in HTF, 0.5% BSA under mineral oil for 4 h (37°C, 90% N₂, 5% O₂, and 5% CO₂) in a BT37GP incubator (Planer). Eggs/embryos/oocytes were washed by serial transfer through 500 µl of HTF/BSA and fixed for imaging (Baibakov et al., 2012). For each experiment using human sperm, initial evaluations were conducted using 10 huZP2^{Rescue} eggs as positive controls to test batches of HTF media and quality of human sperm preparations, which varied.

To assess fertility, females (≥5) from each mouse line were singly co-caged with a fertile female (control, NIH Swiss) and mated (2:1) with a male (NIH Swiss) proven to be fertile. Litters were recorded until the NIH Swiss fertile female gave birth to at least three litters or after 5 mo of mating.

In vivo transcervical insemination

Female mice (8–12 wk old) were hormonally stimulated at 12:00 midnight with 5 IU of pregnant mare serum gonadotrophin (PMSG), and hCG was injected intraperitoneally 48 h later. Epididymal mouse sperm from three males was released into 1.5 ml of HTF, 0.4% BSA, and equilibrated for 1–5 h (37°C, 90% N₂, 5% O₂, and 5% CO₂). 500 µl was used to inseminate each of two female mice. Alternately, commercially obtained human semen (proven to bind and penetrate human ZP2^{Rescue} eggs in vitro) was prepared as described in "Assessment of sperm binding and fertility" and diluted in 500 µl HTF 0.5% BSA that had been equilibrated (≥24 h) at 37°C, 90% N₂, 5% O₂, and 5% CO₂. After capacitation (1.5 h), the contents of two vials of human semen were used to inseminate a single female mouse. At 11:00 p.m. (11 h after hCG injection), females were restrained and inseminated with mouse or human sperm using an NSET device (ParaTechs) connected to a 1-ml syringe, which reached a single uterine horn without anesthesia and surgery (Snell et al., 1944). Females were mated overnight with sterile, vasectomized males, and ~10 h later, eggs were collected from the oviducts of females with copulatory plugs.

In vivo oviduct transfer of sperm

HuZP2^{Rescue} and huZP3^{Rescue} females were stimulated with gonadotrophins and anesthetized. Human sperm were injected into the oviduct, proximal to the swollen ampulla. 2 h later, the mice were euthanized to collect eggs that were fixed and imaged by confocal microscopy (Sato and Kimura, 2001; Tokuhira et al., 2012).

Immunoblot

Ovulated eggs with intact zonae pellucidae from moQuad^(huZP4) and moQuad-moZP2^{trunc} transgenic mice were resolved by SDS-PAGE and probed with monoclonal antibodies to the N and C termini of mouse ZP2 (Burkart et al., 2012).

Peptide bead binding assay

cDNA encoding human (39–154 aa) and mouse (35–149 aa) ZP2 were cloned into pFastBac-HBM TOPO (Invitrogen) downstream of a polyhedron promoter and a 23-aa honeybee melittin signal peptide. Each clone was tagged with 6-histidine at the C terminus to facilitate purification. Using the human cDNA as a backbone, chimeric mouse–human ZP2 clones were generated by DNA synthesis and used to make additional baculovirus expression constructs in which sequences encoding mouse ZP2^{52–83}, ZP2^{85–101}, and ZP2^{103–133} replaced endogenous human sequence. Recombinant peptides were expressed in Hi-Five cells, purified on IMAC Sepharose High Performance beads (GE Healthcare), and assayed on SDS-PAGE (Fig. S2 C) as described previously (Baibakov et al., 2012). Results reflect the mean \pm SEM from at least three independently obtained samples, each containing 20–25 beads. Cloning, expression, and attachment to IMAC beads were performed in the Protein Expression Laboratory of the Advanced Technology Program, SAIC-Frederick, Inc.

Review board compliance

Experiments with normal and transgenic mice were conducted in compliance with the guidelines of the Animal Care and Use Committee of the National Institutes of Health under a Division of Intramural Research, National Institute of Diabetes and Digestive and Kidney Diseases–approved animal study protocol. Human sperm-binding assays were conducted in compliance with an Institutional Review Board approved protocol, and the Office of Human Subject Research determined that federal regulations did not apply to the anonymous human sperm and noninseminated, immature oocytes.

Online supplemental material

Fig. S1 shows the exon maps of transgenic mice and documents their tissue-specific expression. Fig. S2 provides a schematic of transcervical insemination, documents mouse sperm binding to huQuad, huQuad-ZP2^{Null}, and huQuad-ZP2^{Null}; Zp2^{Mo} eggs, and resolves the mouse–human chimeric peptides expressed in baculovirus on SDS-PAGE. Table S1 summarizes the protein composition of the zonae pellucidae of the transgenic mouse lines, and Table S2 summarizes their fertility. Tables S3–S5 list primers used for evaluating transgenes (Table S3), genotyping (Table S4), and RT-PCR (Table S5). Online supplemental material is available at <http://www.jcb.org/cgi/content/full/jcb.201404025/DC1>.

We thank Galina Baibakov for her excellent animal husbandry of the mouse colony necessary for these investigations, and our colleagues for their useful discussions. Noninseminated, immature human oocytes were obtained from Shady Grove Fertility Clinic (Rockville, MD).

This research was supported by the Intramural Research Program of the National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases (ZIA-DK015603).

The authors declare no competing financial interests.

Submitted: 7 April 2014

Accepted: 16 May 2014

References

Baibakov, B., L. Gauthier, P. Talbot, T.L. Rankin, and J. Dean. 2007. Sperm binding to the zona pellucida is not sufficient to induce acrosome exocytosis. *Development*. 134:933–943. <http://dx.doi.org/10.1242/dev.02752>

Baibakov, B., N.A. Boggs, B. Yauger, G. Baibakov, and J. Dean. 2012. Human sperm bind to the N-terminal domain of ZP2 in humanized zonae pellucidae in transgenic mice. *J. Cell Biol.* 197:897–905. <http://dx.doi.org/10.1083/jcb.201203062>

Bauskin, A.R., D.R. Franken, U. Eberspaecher, and P. Donner. 1999. Characterization of human zona pellucida glycoproteins. *Mol. Hum. Reprod.* 5:534–540. <http://dx.doi.org/10.1093/molehr/5.6.534>

Bedford, J.M. 1977. Sperm/egg interaction: the specificity of human spermatozoa. *Anat. Rec.* 188:477–487. <http://dx.doi.org/10.1002/ar.1091880407>

Bleil, J.D., and P.M. Wassarman. 1980. Structure and function of the zona pellucida: identification and characterization of the proteins of the mouse

oocyte's zona pellucida. *Dev. Biol.* 76:185–202. [http://dx.doi.org/10.1016/0012-1606\(80\)90371-1](http://dx.doi.org/10.1016/0012-1606(80)90371-1)

Bleil, J.D., and P.M. Wassarman. 1988. Galactose at the nonreducing terminus of O-linked oligosaccharides of mouse egg zona pellucida glycoprotein ZP3 is essential for the glycoprotein's sperm receptor activity. *Proc. Natl. Acad. Sci. USA*. 85:6778–6782. <http://dx.doi.org/10.1073/pnas.85.18.6778>

Bleil, J.D., C.F. Beall, and P.M. Wassarman. 1981. Mammalian sperm-egg interaction: fertilization of mouse eggs triggers modification of the major zona pellucida glycoprotein, ZP2. *Dev. Biol.* 86:189–197. [http://dx.doi.org/10.1016/0012-1606\(81\)90329-8](http://dx.doi.org/10.1016/0012-1606(81)90329-8)

Boja, E.S., T. Hoodbhoy, H.M. Fales, and J. Dean. 2003. Structural characterization of native mouse zona pellucida proteins using mass spectrometry. *J. Biol. Chem.* 278:34189–34202. <http://dx.doi.org/10.1074/jbc.M304026200>

Bork, P., and C. Sander. 1992. A large domain common to sperm receptors (Zp2 and Zp3) and TGF- β type III receptor. *FEBS Lett.* 300:237–240. [http://dx.doi.org/10.1016/0014-5793\(92\)80853-9](http://dx.doi.org/10.1016/0014-5793(92)80853-9)

Bukovsky, A., S.K. Gupta, P. Bansal, S. Chakravarty, M. Chaudhary, M. Svetlikova, R.S. White, P. Copas, N.B. Upadhyaya, S.E. Van Meter, and M.R. Caudle. 2008. Production of monoclonal antibodies against recombinant human zona pellucida glycoproteins: utility in immunolocalization of respective zona proteins in ovarian follicles. *J. Reprod. Immunol.* 78:102–114. <http://dx.doi.org/10.1016/j.jri.2007.10.004>

Burkart, A.D., B. Xiong, B. Baibakov, M. Jiménez-Movilla, and J. Dean. 2012. Ovastacin, a cortical granule protease, cleaves ZP2 in the zona pellucida to prevent polyspermy. *J. Cell Biol.* 197:37–44. <http://dx.doi.org/10.1083/jcb.201112094>

Chen, J., E.S. Litscher, and P.M. Wassarman. 1998. Inactivation of the mouse sperm receptor, mZP3, by site-directed mutagenesis of individual serine residues located at the combining site for sperm. *Proc. Natl. Acad. Sci. USA*. 95:6193–6197. <http://dx.doi.org/10.1073/pnas.95.11.6193>

East, I.J., and J. Dean. 1984. Monoclonal antibodies as probes of the distribution of ZP-2, the major sulfated glycoprotein of the murine zona pellucida. *J. Cell Biol.* 98:795–800. <http://dx.doi.org/10.1083/jcb.98.3.795>

East, I.J., B.J. Gulyas, and J. Dean. 1985. Monoclonal antibodies to the murine zona pellucida protein with sperm receptor activity: effects on fertilization and early development. *Dev. Biol.* 109:268–273. [http://dx.doi.org/10.1016/0012-1606\(85\)90454-3](http://dx.doi.org/10.1016/0012-1606(85)90454-3)

Florman, H.M., and P.M. Wassarman. 1985. O-linked oligosaccharides of mouse egg ZP3 account for its sperm receptor activity. *Cell*. 41:313–324. [http://dx.doi.org/10.1016/0092-8674\(85\)90084-4](http://dx.doi.org/10.1016/0092-8674(85)90084-4)

Gahlay, G., L. Gauthier, B. Baibakov, O. Epifano, and J. Dean. 2010. Gamete recognition in mice depends on the cleavage status of an egg's zona pellucida protein. *Science*. 329:216–219. <http://dx.doi.org/10.1126/science.1188178>

Ganguly, A., A. Bukovsky, R.K. Sharma, P. Bansal, B. Bhandari, and S.K. Gupta. 2010. In humans, zona pellucida glycoprotein-1 binds to spermatozoa and induces acrosomal exocytosis. *Hum. Reprod.* 25:1643–1656. <http://dx.doi.org/10.1093/humrep/deq105>

Larkin, M.A., G. Blackshields, N.P. Brown, R. Chenna, P.A. McGettigan, H. McWilliam, F. Valentin, I.M. Wallace, A. Wilm, R. Lopez, et al. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics*. 23:2947–2948. <http://dx.doi.org/10.1093/bioinformatics/btm404>

Le Naour, F., E. Rubinstein, C. Jasmin, M. Prenant, and C. Boucheix. 2000. Severely reduced female fertility in CD9-deficient mice. *Science*. 287:319–321. <http://dx.doi.org/10.1126/science.287.5451.319>

Lefèvre, L., S.J. Conner, A. Salpekar, O. Olufowobi, P. Ashton, B. Pavlovic, W. Lenton, M. Afnan, I.A. Brewis, M. Monk, et al. 2004. Four zona pellucida glycoproteins are expressed in the human. *Hum. Reprod.* 19:1580–1586. <http://dx.doi.org/10.1093/humrep/deh301>

Liu, C., E.S. Litscher, and P.M. Wassarman. 1995. Transgenic mice with reduced numbers of functional sperm receptors on their eggs reproduce normally. *Mol. Biol. Cell*. 6:577–585. <http://dx.doi.org/10.1091/mbc.6.5.577>

Liu, C., E.S. Litscher, S. Mortillo, Y. Sakai, R.A. Kinloch, C.L. Stewart, and P.M. Wassarman. 1996. Targeted disruption of the mZP3 gene results in production of eggs lacking a zona pellucida and infertility in female mice. *Proc. Natl. Acad. Sci. USA*. 93:5431–5436. <http://dx.doi.org/10.1073/pnas.93.11.5431>

Liu, P., N.A. Jenkins, and N.G. Copeland. 2003. A highly efficient recombineering-based method for generating conditional knockout mutations. *Genome Res.* 13:476–484. <http://dx.doi.org/10.1101/gr.749203>

Lowe, J.B., and J.D. Marth. 2003. A genetic approach to Mammalian glycan function. *Annu. Rev. Biochem.* 72:643–691. <http://dx.doi.org/10.1146/annurev.biochem.72.121801.161809>

Miller, D.J., M.B. Macek, and B.D. Shur. 1992. Complementarity between sperm surface beta-1,4-galactosyltransferase and egg-coat ZP3 mediates sperm-egg binding. *Nature*. 357:589–593. <http://dx.doi.org/10.1038/357589a0>

Pang, P.C., P.C. Chiu, C.L. Lee, L.Y. Chang, M. Panico, H.R. Morris, S.M. Haslam, K.H. Khoo, G.F. Clark, W.S. Yeung, and A. Dell. 2011. Human sperm binding is mediated by the sialyl-Lewis(x) oligosaccharide

on the zona pellucida. *Science*. 333:1761–1764. <http://dx.doi.org/10.1126/science.1207438>

- Quesada, V., L.M. Sánchez, J. Alvarez, and C. López-Otín. 2004. Identification and characterization of human and mouse ovastacin: a novel metalloproteinase similar to hatching enzymes from arthropods, birds, amphibians, and fish. *J. Biol. Chem.* 279:26627–26634. <http://dx.doi.org/10.1074/jbc.M401588200>
- Rankin, T., M. Familiar, E. Lee, A.M. Ginsberg, N. Dwyer, J. Blanchette-Mackie, J. Drago, H. Westphal, and J. Dean. 1996. Mice homozygous for an insertional mutation in the *Zp3* gene lack a zona pellucida and are infertile. *Development*. 122:2903–2910.
- Rankin, T.L., Z.-B. Tong, P.E. Castle, E. Lee, R. Gore-Langton, L.M. Nelson, and J. Dean. 1998. Human *ZP3* restores fertility in *Zp3* null mice without affecting order-specific sperm binding. *Development*. 125:2415–2424.
- Rankin, T., P. Talbot, E. Lee, and J. Dean. 1999. Abnormal zonae pellucidae in mice lacking *ZP1* result in early embryonic loss. *Development*. 126:3847–3855.
- Rankin, T.L., M. O'Brien, E. Lee, K.E.J.J. Wigglesworth, J. Eppig, and J. Dean. 2001. Defective zonae pellucidae in *Zp2*-null mice disrupt folliculogenesis, fertility and development. *Development*. 128:1119–1126.
- Rankin, T.L., J.S. Coleman, O. Epifano, T. Hoodbhoy, S.G. Turner, P.E. Castle, E. Lee, R. Gore-Langton, and J. Dean. 2003. Fertility and taxon-specific sperm binding persist after replacement of mouse sperm receptors with human homologs. *Dev. Cell*. 5:33–43. [http://dx.doi.org/10.1016/S1534-5807\(03\)00195-3](http://dx.doi.org/10.1016/S1534-5807(03)00195-3)
- Sato, M., and M. Kimura. 2001. Intrabursal transfer of spermatozoa (ITS): a new route for artificial insemination of mice. *Theriogenology*. 55:1881–1890. [http://dx.doi.org/10.1016/S0093-691X\(01\)00529-5](http://dx.doi.org/10.1016/S0093-691X(01)00529-5)
- Shi, S., S.A. Williams, A. Seppo, H. Kurniawan, W. Chen, Z. Ye, J.D. Marth, and P. Stanley. 2004. Inactivation of the *Mgat1* Gene in Oocytes Impairs Oogenesis, but Embryos Lacking Complex and Hybrid N-Glycans Develop and Implant. *Mol. Cell. Biol.* 24:9920–9929. <http://dx.doi.org/10.1128/MCB.24.22.9920-9929.2004>
- Snell, G.D., K.P. Hummel, and W.H. Abelman. 1944. A technique for the artificial insemination of mice. *Anat. Rec.* 90:243–253. <http://dx.doi.org/10.1002/ar.1090900309>
- Spargo, S.C., and R.M. Hope. 2003. Evolution and nomenclature of the zona pellucida gene family. *Biol. Reprod.* 68:358–362. <http://dx.doi.org/10.1095/biolreprod.102.008086>
- Thall, A.D., P. Malý, and J.B. Lowe. 1995. Oocyte Gal α 1,3Gal Epitopes Implicated in Sperm Adhesion to the Zona Pellucida Glycoprotein ZP3 Are Not Required for Fertilization in the Mouse. *J. Biol. Chem.* 270:21437–21440.
- Tian, J., H. Gong, and W.J. Lennarz. 1999. *Xenopus laevis* sperm receptor gp69/64 glycoprotein is a homolog of the mammalian sperm receptor ZP2. *Proc. Natl. Acad. Sci. USA*. 96:829–834. <http://dx.doi.org/10.1073/pnas.96.3.829>
- Tokuhiro, K., M. Ikawa, A.M. Benham, and M. Okabe. 2012. Protein disulfide isomerase homolog PDILT is required for quality control of sperm membrane protein ADAM3 and male fertility [corrected]. *Proc. Natl. Acad. Sci. USA*. 109:3850–3855. <http://dx.doi.org/10.1073/pnas.1117963109>
- Warming, S., N. Costantino, D.L. Court, N.A. Jenkins, and N.G. Copeland. 2005. Simple and highly efficient BAC recombineering using galK selection. *Nucleic Acids Res.* 33:e36. <http://dx.doi.org/10.1093/nar/gni035>
- Williams, S.A., L. Xia, R.D. Cummings, R.P. McEver, and P. Stanley. 2007. Fertilization in mouse does not require terminal galactose or N-acetylglucosamine on the zona pellucida glycans. *J. Cell Sci.* 120:1341–1349. <http://dx.doi.org/10.1242/jcs.004291>
- Yauger, B., N.A. Boggs, and J. Dean. 2011. Human *ZP4* is not sufficient for taxon-specific sperm recognition of the zona pellucida in transgenic mice. *Reproduction*. 141:313–319. <http://dx.doi.org/10.1530/REP-10-0241>
- Zhao, M., L. Gold, A.M. Ginsberg, L.-F. Liang, and J. Dean. 2002. Conserved furin cleavage site not essential for secretion and integration of *ZP3* into the extracellular egg coat of transgenic mice. *Mol. Cell. Biol.* 22:3111–3120. <http://dx.doi.org/10.1128/MCB.22.9.3111-3120.2002>